 Your sex partner just told you that they have been diagnosed with trichomoniasis (trich). This infection is spread through vaginal sex. You may be infected and have no symptoms. If left untreated, it can increase your risk of getting passing HIV. Get Tested & Treated Take this card to your doctor, clinic, or LA County STD Clinic today. For more information, clinic referrals, and free condoms, call the STD Hotline 1+800-758-0880 or visit GetProtectedLA.com.	 Your sex partner just told you that they have been diagnosed with trichomoniasis (trich). This infection is spread through vaginal sex. You may be infected and have no symptoms. If left untreated, it can increase your risk of getting passing HIV. Get Tested & Treated Take this card to your doctor, clinic, or LA County STD Clinic today. For more information, clinic referrals, and free condoms, call the STD Hotline 1-800-758-0880 or visit GetProtectedLA.com. 		
		Your sex partner just told you that they have been diagnosed with <u>trichomoniasis (trich)</u> . This infection is spread through vaginal sex. 	Your sex partner just told you that they have been diagnosed with <u>trichomoniasis (trich)</u> . This infection is spread through vaginal sex.
		 You may be infected and have no symptoms. If left untreated, it can increase your risk of getting passing HIV. Get Tested & Treated Take this card to your doctor, clinic, 	 You may be infected and have no symptoms. If left untreated, it can increase your risk of getting passing HIV. Get Tested & Treated Take this card to your doctor, clinic,
or LA County STD Clinic today. For more information, clinic referrals, and free condoms, call the STD Hotline -800-758-0880 or visit GetProtectedLA.com.	or LA County STD Clinic today. For more information, clinic referrals, and free condoms, call the STD Hotline 1-800-758-0880 or visit <u>GetProtectedLA.com</u> .		
 Your sex partner just told you that they have been diagnosed with trichomoniasis (trich). This infection is spread through vaginal sex. You may be infected and have no symptoms. If left untreated, it can increase your risk of getting passing HIV. 	 Your sex partner just told you that they have been diagnosed with <u>trichomoniasis (trich)</u>. This infection is spread through vaginal sex. You may be infected and have no symptoms. If left untreated, it can increase your risk of getting passing HIV. 		
Get Tested & Treated Take this card to your doctor, clinic, or LA County STD Clinic today.	Get Tested & Treated Take this card to your doctor, clinic, or LA County STD Clinic today.		
or more information, clinic referrals, and free condoms, call the STD Hotline -800-758-0880 or visit GetProtectedLA.com.	For more information, clinic referrals, and free condoms, call the STD Hotline 1-800-758-0880 or visit <u>GetProtectedLA.com</u> .		
Your sex partner just told you that they have been liagnosed with <u>trichomoniasis (trich)</u> . • This infection is spread through vaginal sex. • You may be infected and have no symptoms.	 Your sex partner just told you that they have been diagnosed with <u>trichomoniasis (trich)</u>. This infection is spread through vaginal sex. You may be infected and have no symptoms. 		
 If left untreated, it can increase your risk of getting passing HIV. Get Tested & Treated Take this card to your doctor, clinic, or LA County STD Clinic today. 	 If left untreated, it can increase your risk of getting passing HIV. Get Tested & Treated Take this card to your doctor, clinic, or LA County STD Clinic today. 		
or more information, clinic referrals, and free condoms, call the STD Hotline -800-758-0880 or visit GetProtectedLA.com.	For more information, clinic referrals, and free condoms, call the STD Hotline 1-800-758-0880 or visit GetProtectedLA.com.		
Your sex partner just told you that they have been diagnosed with <u>trichomoniasis (trich)</u> .	Your sex partner just told you that they have been diagnosed with trichomoniasis (trich).		
 This infection is spread through vaginal sex. You may be infected and have no symptoms. If bett untracted, it can increase your risk of actting pageing LIV. 	 This infection is spread through vaginal sex. You may be infected and have no symptoms. 		
 If left untreated, it can increase your risk of getting passing HIV. Get Tested & Treated Take this card to your doctor, clinic, 	 If left untreated, it can increase your risk of getting passing HIV. Get Tested & Treated Take this card to your doctor, clinic, 		
or LA County STD Clinic today.	or LA County STD Clinic today.		

Get Tested & Treated Take this card to your doctor, clinic, or LA County STD Clinic today.

For more information, clinic referrals, and free condoms, call the STD Hotline 1-800-758-0880 or visit <u>GetProtectedLA.com</u>.

For more information, clinic referrals, and free condoms, call the STD Hotline 1-800-758-0880 or visit <u>GetProtectedLA.com</u>.

or LA County STD Clinic today.

For doctor or nurse:

The person with this card has been exposed to trichomoniasis. The CDC recommends sexual contacts to chlamydia be treated presumptively with:

> Metronidazole 2 g orally in single dose - OR -Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacountv.gov/dhsp/InfoForProviders.htm

Rublic Health Pavisad 03/2022

KPublic Health

EX Public Health

Revised: 03/2022

Revised: 03/2022

For doctor or nurse:

The person with this card has been exposed to trichomoniasis. The CDC recommends sexual contacts to chlamydia be treated presumptively with:

> Metronidazole 2 g orally in single dose - OR -Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacountv.gov/dhsp/InfoForProviders.htm

For doctor or nurse:

The person with this card has been exposed to trichomoniasis. The CDC recommends sexual contacts to chlamydia be treated presumptively with:

> Metronidazole 2 g orally in single dose - OR -

Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacounty.gov/dhsp/InfoForProviders.htm

For doctor or nurse:

The person with this card has been exposed to trichomoniasis. The CDC recommends sexual contacts to chlamydia be treated presumptively with:

Metronidazole 2 g orally in single dose - OR Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacounty.gov/dhsp/InfoForProviders.htm

Expublic Health Revised: 03/2022

For doctor or nurse:

The person with this card has been exposed to trichomoniasis. The CDC recommends sexual contacts to chlamydia be treated presumptively with:

> Metronidazole 2 g orally in single dose - OR -

Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacounty.gov/dhsp/InfoForProviders.htm

EX Public Health Revised: 03/2022

For doctor or nurse:

The person with this card has been exposed to trichomoniasis. The CDC recommends sexual contacts to chlamydia be treated presumptively with:




Metronidazole 2 g orally in single dose - OR -

Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, **K**Public Health visit www.publichealth.lacountv.gov/dhsp/InfoForProviders.htm





Revised: 03/2022

The person with this card has been exposed to trichomoniasis The CDC recommends sexual contacts to chlamydia be treated presumptively with:

> Metronidazole 2 g orally in single dose - OR -

Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacounty.gov/dhsp/InfoForProviders.htm



For doctor or nurse:

The person with this card has been exposed to trichomoniasis. The CDC recommends sexual contacts to chlamydia be treated presumptively with:

> Metronidazole 2 g orally in single dose - OR -

Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacounty.gov/dhsp/InfoForProviders.htm



For doctor or nurse:

The person with this card has been exposed to trichomoniasis, The CDC recommends sexual contacts to chlamydia be treated presumptively with:

> Metronidazole 2 g orally in single dose - OR -

Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacounty.gov/dhsp/InfoForProviders.htm



For doctor or nurse:

The person with this card has been exposed to trichomoniasis. The CDC recommends sexual contacts to chlamydia be treated presumptively with:

> Metronidazole 2 g orally in single dose - OR -

Metronidazole 500 mg orally twice a day for 7 days

Testing for chlamydia, gonorrhea, syphilis, and HIV is also recommended.

For completed CDC treatment guidelines and other STD information, visit www.publichealth.lacounty.gov/dhsp/InfoForProviders.htm





