### SYPHILIS SYMPTOMS

- The first symptom is a painless sore that may appear inside or near the mouth, genitals or anal areas.
- A second symptom may be a rash that appears on the palms of hands, bottom of feet, or on the body between the arm pits and upper thighs.
- Syphilis symptoms may appear and disappear.
- To cure syphilis medical testing and treatment are required.



### **TEST TODAY**

- If experiencing signs or symptoms
- If being tested for other STDs
- If sexually active (testing recommended every 6-12 months)
- If engaging in unprotected sex (testing recommended every 6 months)
- If you have multiple sex partners (testing recommended every 6 months)
- If you are a man having sex with other men (testing recommended every 3-6 months)
- If you plan to get pregnant in the future

# That sore might be SYPHILIS Syphilis and other STDs are on the rise in LA County GET FACTS. GET TESTED. GET CURED.

Free & confidential testing & information: 1-800-758-0880 • www.publichealth.lacounty.gov/dhsp



## It's their health too SYPHILIS

and other STDs can harm your baby GET FACTS. GET TESTED. GET CURED.

Free & confidential testing & information: 1-800-758-0880 • www.publichealth.lacounty.gov/dhsp



### **SYPHILIS IN WOMEN**

- Syphilis in women is increasing in Los Angeles County
- If a woman is infected with syphilis while pregnant, it can lead to birth defects, miscarriage, premature birth or stillbirth
- Good News: Syphilis can be treated and cured even during pregnancy with a shot of antibiotics

#### IF YOU ARE PREGNANT:

- Prenatal care is very important to keep you and your baby healthy
- You should be tested for syphilis 3 times during your pregnancy: during the 1st trimester, early in the 3rd trimester (between 28-32 weeks) and at delivery
- If you don't have a regular doctor, call the free STD hotline (800) 758-0880 to find out where to get tested and treated for free